



Certificate of Analysis

Nelson Honey & Marketing (NZ) Ltd
 276 State Highway 6, RD 2, Motupiko
 Nelson 7072
 Attention: Terena Roeske
 Phone: 03 522 4413
 Email: terena@nelsonhoney.com

Lab Reference: 17-15584
 Submitted by:
 Date Received: 29/06/2017
 Date Completed: 3/07/2017
 Order Number:
 Reference:

Report Comments

Samples were received by Analytica Laboratories in acceptable condition unless otherwise noted on this report.

Results Summary

3in1 Honey Analysis

Laboratory ID	Sample ID	Dihydroxyacetone DHA	Methylglyoxal MG	Non-peroxide Activity NPA*	Hydroxymethylfurfural HMF
		<i>Units</i> mg/kg	mg/kg	%w/v phenol eq.	mg/kg
		<i>Reporting Limit</i> 10	4	0.8	1
17-15584-1	230617MH	300	151	7.2	27

3in1 Honey Analysis Approver:

Jacob Jaine, Ph.D.
 Senior Technologist

Method Summary

3in1 Determination of Dihydroxyacetone (DHA), Methylglyoxal (MG) and Hydroxymethylfurfural (HMF) by aqueous extraction, derivatisation, and UPLC analysis.

NPA Non-Peroxide Activity (NPA) values are not directly measured by the laboratory, but are calculated from the measured methylglyoxal concentration in the honey according to the requirements of the client. The calculation is based on published data^(†) comparing the NPA and methylglyoxal concentration measured in a range of honey samples. These calculated values are not accredited by IANZ and do not imply that the honey is or is not manuka honey. NPA values less than 5 are an estimate based on extrapolation of the relationship between methylglyoxal and NPA

(†) Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. And, Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey" [Carbohydr. Res. 343 (2008) 651]. Carbohydrate Research 344 (2009) 2609. C. J. Adams, et al.